



# Technical Update 1

## Manganese Deficiency & Marsh Spot

January 2025

A deficiency or unavailability of the trace element manganese can cause a reduction of yield in both vining and combining peas and may result in the appearance of marsh spot, a disorder that affects the seed, and reduces produce quality for human consumption and affects germination of seedlings.

### OCCURRENCE AND SYMPTOMS

The disorder is commonly associated with organic soils, although crops on other soil types can also be affected. Peas grown on soils with a pH greater than 6.8 are more susceptible to deficiency, especially in wet or compacted conditions, when manganese becomes unavailable to the plants and is locked-up in the alkaline soil. Over liming must be avoided. Heavily trafficked areas, such as wheelings, cause compaction, and symptoms of manganese deficiency can often be seen as pale coloured lines running through a crop.

Manganese deficiency causes a yellowing around the leaf edges and between the veins. Entire leaves may become yellow in time, but the veins and the areas of tissue adjoining them are the last to lose their green colour, giving a somewhat striped appearance to the leaf.

Affected plants are likely to produce peas which have a brown or granular necrotic area in the centre of the cotyledons and this condition is known as marsh spot. Such produce is unsuitable for processing due to the physical blemish, or for use as seed, as the spot affects the plumule and gives rise to abnormally developed seedlings.

### TREATMENTS

Treat crops which are:

- Growing on land known to have a history of manganese deficiency.
- Growing on land of pH 6.8 or more as a routine, spraying at the 1st pod stage and again 10-14 days later.



In addition:

- During wet weather flowering may be prolonged and a third application may be beneficial.
- Crops in which symptoms are seen should be sprayed immediately and treatment repeated at the 1st pod stage and about 10 days later still.

Treat with manganese sulphate (32% manganese) at 5 kg/ha in a high volume of water with an added wetter. Trials have shown that other products containing a high percentage of manganese are also effective. However, sprays with chelated forms (6% manganese) do not contain enough manganese to give adequate control.

### COMBINED TREATMENTS

Sometimes, treatment against manganese deficiency coincides with treatment against pea moth or pea aphid or with a fungicide. Certain insecticides and fungicides can be mixed, but before doing so, the manufacturers of the products must be consulted.

The information in this publication must not be reproduced without the express written permission from the PGRO.

Information disseminated by the Processors & Growers Research Organisation is given after the exercise of all possible care in compilation, preparation and issue, but is provided without liability in its application or use.

## Processors & Growers Research Organisation

The Research Station, Great North Road, Thornhaugh, Cambridgeshire PE8 6HJ, UK  
**Chief Executive:** Roger Vickers **Tel:** +44 (0)1780 782585 **email:** info@pgro.org **www:** pgro.org  
Company Limited by Guarantee - Registered in London No. 567232. Registered Charity No 284077. VAT Reg No 281 7742 37GB