



Technical Update 06

Black bean aphid

January 2024

Dense colonies of black aphids can develop at any time on the upper part of faba bean stems from the onset of flowering. The colonies can consist of hundreds of aphids which gradually extend downwards onto the leaves and developing pods. At first, individual plants may be infested and later the aphids spread into localised patches of infested crop.



Black bean aphid colony

DAMAGE

Damage to bean crops occurs mainly as the direct effect of aphids feeding. Plants fail to develop, and pods do not fill normally. The honeydew produced during feeding attracts secondary moulds as well as *Botrytis* spp. which cause chocolate spot. In addition, black bean aphids transmit viruses such as bean leaf roll virus.

DESCRIPTION AND LIFE CYCLE

Individual aphids are black, 1-2 mm in size, with very small white spots on the upper surface of the body. Winged aphids migrate to beans in early summer from their over-wintering hosts, which include the spindle bush (*Euonymus europaeus*). Each adult is viviparous and can produce several wingless nymphs each day. Colonies develop rapidly, particularly in warm, humid conditions.


VIRUSES

Bean leaf roll virus (BLRV)

BLRV is aphid-transmitted and both the black bean aphid and pea aphid are vectors. Infection is usually more obvious where aphids infest the crop before flowering. The pea aphid is more likely to move into beans currently. Control of early invading aphids is essential to prevent virus infection.

Pea enation mosaic virus (PEMV)

The upper leaves of the plants may become pointed and crinkled. Vein clearing can develop, and translucent spotting may appear. Individual plants or small groups of plants can show symptoms from late flowering onwards. Pea aphids are the most common vectors. Control of aphids at early flowering is likely to prevent serious infection with PEMV.



Bean yellow mosaic virus

Bean yellow mosaic virus (BYMV)

BYMV develops in beans at any time before flowering. Leaves are crinkled and may become pointed. Vein clearing can develop, and the plant is slightly stunted. It is similar to PEMV but without the translucent spotting and streaking on the leaf surface. The virus is aphid transmitted and common in field beans. Infection may be on individual plants or groups of plants and, depending on severity, yield can be severely reduced. Pea aphid is the principal vector, although black bean aphid can also transmit the virus. Early control of aphids in flowering bean crops is the most effective means of preventing virus infection.

CONTROL IN THE FIELD

Black bean aphids are relatively easy to control using aphicide sprays, but because the beans are often flowering, the choice of aphicides is limited to those which will not harm pollinating insects. Colonies first develop on headlands, but where a general infestation of 10% of plants are colonised, spraying should be carried out as soon as possible. Applications made at early flowering or when 5% of plants are infested reduce infections of aphid-transmitted viruses. The Rothamsted Insect Survey (<https://insectsurvey.com/aphid-bulletin>) provides information about aphid flights throughout the season and is a useful tool to help predict activity in crops.

The information in this publication must not be reproduced without the express written permission from the PGRO.

Information disseminated by the Processors & Growers Research Organisation is given after the exercise of all possible care in compilation, preparation and issue, but is provided without liability in its application or use.

Processors & Growers Research Organisation

The Research Station, Great North Road, Thornhaugh, Cambridgeshire PE8 6HJ, UK
Chief Executive: Roger Vickers **Tel:** +44 (0)1780 782585 **email:** info@pgro.org www.pgro.org
Company Limited by Guarantee - Registered in London No. 567232. Registered Charity No 284077. VAT Reg No 281 7742 37GB